

## **Caspialosa caspia tanaica**

2010-08-30 22:09:24 by admin

**Caspialosa caspia tanaica** Family : Clupeidae Genus : Caspialosa Species : caspia tanaica **Accepted scientific name:** *Alosa tanaica* (Grimm, 1901) (accepted name) **Synonyms:** *Alosa bulgarica* Drensky, 1934 (synonym) *Alosa caspia nordmanni* Antipa, 1904 (synonym) *Alosa caspia tanaica* (Grimm, 1901) (synonym) *Alosa caspia tanaica asovi* Pavlov, 1959 (ambiguous synonym) *Alosa nordmanni* Antipa, 1904 (synonym) *Caspialosa caspia palaeostomi* Sadowsky, 1934 (synonym) *Caspialosa caspia tanaica* (Grimm, 1901) (synonym) *Caspialosa palaeostomica* Sadowsky, 1934 (synonym) *Caspialosa tanaica etemi* Battalgil, 1941 (ambiguous synonym) *Caspialosa tanaica palaeostomi* Sadowsky, 1934 (synonym) *Caspiolosa tanaica etemi* Battalgil, 1941 (ambiguous synonym) *Clupea tanaica* Grimm, 1901 (synonym) **Size / Weight / Age** Max length : 20.0 cm TL male/unsexed; (Ref. 593); max. published weight: 59.0 g (Ref. 593) **Environment** Pelagic-neritic; anadromous (Ref. 51243); freshwater; brackish; marine **Climate / Range** Temperate; 49°N - 36°N, 27°E - 54°E

**Caspialosa caspia tanaica** Family : Clupeidae Genus : Caspialosa Species : caspia tanaica **Accepted scientific name:** *Alosa tanaica* (Grimm, 1901) (accepted name) **Synonyms:** *Alosa bulgarica* Drensky, 1934 (synonym) *Alosa caspia nordmanni* Antipa, 1904 (synonym) *Alosa caspia tanaica* (Grimm, 1901) (synonym) *Alosa caspia tanaica asovi* Pavlov, 1959 (ambiguous synonym) *Alosa nordmanni* Antipa, 1904 (synonym) *Caspialosa caspia palaeostomi* Sadowsky, 1934 (synonym) *Caspialosa caspia tanaica* (Grimm, 1901) (synonym) *Caspialosa palaeostomica* Sadowsky, 1934 (synonym) *Caspialosa tanaica etemi* Battalgil, 1941 (ambiguous synonym) *Caspialosa tanaica palaeostomi* Sadowsky, 1934 (synonym) *Caspiolosa tanaica etemi* Battalgil, 1941 (ambiguous synonym) *Clupea tanaica* Grimm, 1901 (synonym) **Size / Weight / Age** Max length : 20.0 cm TL male/unsexed; (Ref. 593); max. published weight: 59.0 g (Ref. 593) **Environment** Pelagic-neritic; anadromous (Ref. 51243); freshwater; brackish; marine **Climate / Range** Temperate; 49°N - 36°N, 27°E - 54°E

**Distribution** Black Sea, Sea of Marmara and Sea of Azov. **Biology** Migrates for spawning to the lower reaches of the Don in April. At the end of April, it enters the delta of the Kuban and spawns there in June. In winter, it is found along the Caucasian coast as far south as Batumi. It migrates from the Sea of Azov earlier than the other herrings in autumn and returns there in mid-April. Eggs are bathypelagic or sink to bottom **Common names: Common name Language**

**Country** Sled tanajský Czech Czech Rep Azov shad English Former USSR Paeostom shad English Former USSR Rizeafca Rumanian Romania &#1072;&#1079;&#1086;&#1074;&#1089;&#1082;&#1072;&#1103;&#1089;&#1077;&#1083;&#1100;&#1076;&#1100; Russian Russian Fed Palaeostomskii puzanok Russian Former USSR Azovskii puzanok Russian Former USSR Ringa baligi Turkish Turkey